



Nuclear histochemistry: Its history in fifty volumes

M.G. Manfredi Romanini

Dipartimento di Biologia Animale, University of Pavia, Italy

The fiftieth anniversary of the double helix was marked two years ago. The elucidation of the structure of DNA by Watson and Crick can rightfully be considered to be "the" milestone in modern biology because it has conditioned the explosive growth in our understanding of the molecular mechanisms regulating the structural and functional organization of single cells and multicellular organisms.

It is interesting to note how early on histochemical research reflected an awareness of the key role of DNA in the expression of hereditary characters. Ever since Boivin et al. (1948) and Vendrely and Vendrely (1948) established this concept on biochemical bases, there began to appear over time other papers (e.g. Ris and Mirsky, 1949; Swift, 1950; Leuchtemberger et al., 1951 or the largely comparative review article by Vialli, 1957) that established by quantitative cytochemical means that the content of DNA was an important evolutionary parameter in comparing animal and plant taxa. In fact, during those years, the quality of research had also undergone major changes, moving from qualitative analysis *in situ* to quantitative cytochemical measurements, thanks to the introduction of the more sensitive morphophotometric assays, with the first two-way or more-way hand-made photometers coming from Belgium (Lison, 1950a,b, who was the author of the first technical text on quantitative histochemistry: Lison, 1960), Italy (Vialli and Perugini, 1954), and later England (Deeley, 1955). In these studies, it becomes clear that the relative nuclear DNA content is not only a species-specific constant (i.e. found in all cells and all tissues of every individual of a given species) but assumes the significance of an evolutionary parameter because its changes are interpreted to reflect chromosomal or gene mutations, and thus, genetic changes.

The principle that the nuclear DNA content is a constant was in itself a great discovery from the viewpoint of histochemistry because relative quantitative data are used (the intensity of the Feulgen

reaction product in the nucleus relative to the area of the nucleus), relating a chemical colorimetric datum with a morphological one (i.e. the measurement of the size of the nucleus). It was both easy and at the same time biologically correct to use *in situ* techniques to determine the function of single cells, correlating histochemical findings with morphological measurements.

On this basis, in the second half of the last century, with the introduction of fluorescent dyes which permitted the determination of DNA with greater sensitivity, not only quantitative changes in nuclear DNA during the life cycle of the cell, but also alterations in chromatin structure and organization related to the cell metabolic changes in response to different environmental stimuli, could be described. This approach was enormously successful in the analysis of cancerous tissue and is now routinely used in cytodiagnosis.

This copy of the European Journal of Histochemistry is the first issue of volume 50. Beginning as the Rivista di Istochimica Normale e Patologica (Journal of Normal and Pathological Histochemistry), the European Journal of Histochemistry has contributed with its publications to codify the science of histochemistry and has certainly had its impact on the birth and development of the histochemistry of the cell nucleus.

Starting from the front page of the first volume (Figure 1), in fact, the roster of Editors, among them Maffo Vialli, expresses how transversal is the subject matter covered by histochemistry, e.g. basic biology, human and comparative anatomy, and especially microscopic anatomy and pathological and clinical histology. And, at the same time the importance not only of histological methods, but also of physical, physico-chemical, and biochemical methods applied *in situ* are stressed.

The Indices of the first issues purposely listed purely technical papers and more biological articles in which pathological tissues or the changes in normal tissue following pathological stimuli were



Figure 1. Front page of the first issue of the journal *Rivista di Istochimica Normale e Patologica*, 1954.

described. In fact, as early as the second issue, there has appeared an account of a meeting dedicated to histochemistry with particular reference to tumors with some important presentations such as that by Vialli on *histochemical and morphological problems*, by the anatomic-pathologist Sirtori on the use of histochemistry in *the study and diagnosis of tumors* and by the clinician Villa on *nucleic acids in leukemic cells* (Ceriotti, 1954)

In these 50 volumes, in which the history of the European Journal of Histochemistry is written, qualitative and quantitative histochemistry of the nucleus has been an ample and constant factor. More than 230 articles have treated this topic both from basic bio-cellular aspects and from the

implied applications to human pathology. This attitude has become especially apparent during the last few years, when several papers have been published which deserve high interest from the viewpoint of both the improvement of methodology (Mazzini *et al.*, 2003; Bottiroli *et al.*, 2004; Cremer *et al.*, 2004; Pederson, 2004; Marziliano *et al.*, 2005; Sheval *et al.*, 2005) and the application to clinical research (Alexandrakis *et al.*, 2005; Fukuda *et al.*, 2005; Hirose *et al.*, 2005; Soldani *et al.*, 2005).

The Maffo Vialli International Award for Histochemistry, established by the Italian Society Histochemistry and the European Journal of Histochemistry, now in its fifth edition, has previously been assigned to three outstanding scientists

for their distinguished research on the biology of the nucleus: Mels van der Ploeg, in 1999 (Van der Ploeg, 2000), Stan Fakan, in 2003 (Fakan, 2004), and Thomas Cremer, in 2005. Their work demonstrates how histochemical studies cannot but result from the articulated integration of enzymo-histochemical, immunohistochemical and *in situ* hybridization data, more often than not associated with three dimensional computer aided reconstruction of the functional distribution of a particular chemical activity.

Nuclear histochemistry is now therefore a real molecular biology *in situ* applied to research on dynamic processes in the nucleus; and this renders the microscopic and histochemical approach absolutely irreplaceable for the progress in our understanding of cell biology.

Maria Gabriella Manfredi Romanini
Dipartimento di Biologia Animale
Piazza Botta 10, 27100 Pavia, Italy
E-mail: mgmaro@tin.it

References

- Alexandrakis M.G., Passam F.H., Dambaki K., Nalbanti F., Stathopoulos E.S., Kyriakou D.S. The assessment of proliferating cell nuclear antigen immunostaining in myelodysplastic syndromes and its prognostic significance. *Eur J Histochem* 2005; 49: 27-32.
- Boivin A, Vendrely R, Vendrely C. L'acide désoxyribonucléique du noyau cellulaire, dépositaire des caractères héréditaires: atguments diordre analytique. *Compt Rend Acad Sc* 1948; 226:1061-62.
- Bottiroli G, Croce AC, Bottone MG, Vaccino S, Pellicciari C. G0-G1 cell cycle phase transition as revealed by fluorescence resonance energy transfer: analysis of human fibroblast chromatin. *Eur J Histochem* 2004;48:37-48.
- Ceriotti G. Resoconto di una giornata dedicata ai "Problemi di Istochimica con particolare riguardo ai tumori". *Riv Istochim Norm Pat* 1954, 1:239-43.
- Cremer M, Zinner R, Stein S, Albiez H, Wagler B, Cremer C, Cremer T. Three dimensional analysis of histone methylation patterns in normal and tumor cell nuclei. *Eur J Histochem* 2004;48:15-28.
- Deeley EM. An integrating microdensitometer for biological cells. *J Sci Instrum* 1955; 32:263-67.
- Fakan S. Ultrastructural cytochemical analyses of nuclear functional architecture. *Eur J Histochem* 2004;48:5-14.
- Fukuda M, Sun A. The DNA-instability test as a specific marker of malignancy and its application to detect cancer clones in borderline malignancy. *Eur J Histochem* 2005; 49: 11-26.
- Hirose M, Sun A, Okubo T, Noriki S, Imamura Y, Fukuda M. Detection of non-papillary, non-invasive transitional cell G1 carcinoma as revealed by increased DNA instability and other cancer markers. *Eur J Histochem* 2005; 49:199-210.
- Leuchtenberger C, Vendrely R, Vendrely C. A Comparison of the Content of Desoxyribonucleic Acid (DNA) in Isolated Animal Nuclei by Cytochemical and Chemical Methods *PNAS USA* 1951; 37:33-8.
- Lison L. Etude et realisation d'un photometre a l'usage histologique. *Acta Anat* 1950a;10:333-47.
- Lison L. Progres recents de l'histochemie quantitative. *Ann Soc R Sci Med Nat Brux.* 1950b;3:154-68.
- Lison L. Histochemie et cytochimie animales, principes et méthodes. Paris, Gauthier-Villars, 1960.
- Marziliano N., Arbustini E., Rossi de Gasperis M, Crovella S. Detection of Epstein Barr virus in formalin-fixed paraffin tissues by fluorescent direct in situ PCR. *Eur J Histochem* 2005; 49:309-312.
- Mazzini G, Ferrari C, Erba E. Dual excitation multi- fluorescence flow cytometry for detailed analyses of viability and apoptotic cell transition. *Eur J Histochem* 2003;47:289-98.
- Pasteels J, Lison L. Recherches histophotometriques sur la teneur en acide desoxyribonucleique au cours de mitoses somatiques. *Arch Biol* 1950;61:445-74.
- Pederson T. The molecular cytology of gene expression: fluorescent RNA as both a stain and tracer in vivo. *Eur J Histochem* 2004;48:57-64.
- Ris H, Mirsky AE. Quantitative cytochemical determination of desoxyribonucleic acid with the Feulgen nucleal reaction. *J Gen Physiol* 1949; 33:125-46.
- Sheval EV, Polzikov MA, Olson MOJ, Zatssepina OV. A higher concentration of an antigen within the nucleolus may prevent its proper recognition by specific antibodies. *Eur J Histochem* 2005;49:117-24.
- Soldani C., Scovassi A.I., Canosi U., Bramucci E., Ardissino D., Arbustini E. Multicolor fluorescence technique to detect apoptotic cells in advanced coronary atherosclerotic plaques. *Eur J Histochem* 2005; 49:47-52.
- Swift H. The constancy of desoxyribose nucleic acid in plant nuclei. *PNAS USA* 1950; 36:643-54.
- Van der Ploeg M. Cytochemical nucleic acid research during the twentieth century. *Eur J Histochem* 2000;44:7-42.
- Vendrely R, Vendrely, C. La teneur du noyau cellulaire en acide desoxyribonucléique à travers les organes, les individus et les espèces animales. *Experientia* 1948; 4 :434-6.
- Vialli M, Perugini S. Due nuovi modelli di apparecchiature istofotometriche. *Riv Istochim Norm Pat* 1954, 1:149-70.
- Vialli M. Volume et contenu en ADN pour noyau. *Exp Cell Res* 1957; 13:284-93.