

DOI: [10.4081/ejh.2018.2942](https://doi.org/10.4081/ejh.2018.2942)

Expression of acetylated tubulin in the postnatal developing mouse cochlea

Wen Jing Liu,^{1,2} Chuan Xi Wang,¹ Hao Yu,³ Shao Feng Liu,¹ Jun Yang²

¹Department of Otorhinolaryngology-Head and Neck Surgery, Yijishan Hospital, Wannan Medical College, Wuhu


²Department of Otorhinolaryngology-Head and Neck Surgery, Xinhua Hospital, Shanghai Jiaotong University School of Medicine

³Anhui Province Key Laboratory of Biological Macro-Molecules Reserch, Wannan Medical College, Wuhu, China

Correspondence: Jun Yang, Department of Otorhinolaryngology-Head & Neck Surgery, Xinhua Hospital, Shanghai Jiaotong University School of Medicine, Shanghai Jiaotong University Ear Institute, Shanghai Key Laboratory of Translational Medicine on Ear and Nose diseases (No. 14DZ2260300), Shanghai 200092, China. Tel. +86.21.25078532 - Fax: +86.21.65156489. E-mail: yangjun@xinquamed.com

Shao Feng Liu, Department of Otorhinolaryngology-Head and Neck Surgery, Yijishan Hospital, Wannan Medical College, Wuhu, China. E-mail: liusf_cn@163.com

Key words: Acetylated tubulin; immunofluorescence; expression; mouse; cochlea; development.



Supplementary Figure 1.

Detail of P1, P8, P10, P14 and P30 mouse organ of Corti double-labeled by tubulin acetylation (red), myosin VIIa (green) under confocal microscopy between P1 and P30. Micrographs presented in figures were overlaid with (A-C, G-I, J-L) and without (D-F, M-O) bright-field.