

DOI: [10.4081/ejh.2018.2956](https://doi.org/10.4081/ejh.2018.2956)

Interaction between sphingosine kinase/sphingosine 1 phosphate and transforming growth factor- β /Smads pathways in experimental intestinal fibrosis. An *in vivo* immunohistochemical study

Roberta Sferra,^{1§} Simona Pompili,^{1§} Luca Ventura,² Caroline Dubuquoy,³ Silvia Speca,⁴
Eugenio Gaudio,⁵ Giovanni Latella,^{6§} Antonella Vetuschì^{1§}

¹Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, Italy

²Pathology Division, San Salvatore Hospital, L'Aquila, Italy

³Intestinal Biotech Development, Lille, France

⁴U995-LIRIC-Lille Inflammation Research International Center, University of Lille, France

⁵Department of Anatomical, Histological, Forensic Medicine and Orthopedic Sciences, Sapienza University of Rome, Italy


⁶Department of Life, Health and Environmental Sciences, Gastroenterology Unit, University of L'Aquila, Italy

§These authors contributed equally to this work.


Correspondence: Roberta Sferra, Department of Biotechnological and Applied Clinical Sciences, Human Anatomy, University of L'Aquila, Via Vetoio, Coppito 2, 67100, L'Aquila, Italy. Tel.: +39.0862.433574, 433504 - Fax: +39.0862.433523. E-mail: roberta.sferra@univaq.it

Key words: Intestinal fibrosis, epithelial to-mesenchymal transition, immunohistochemistry, TGF β /Smads, sphingosine 1-phosphate.


p-mTOR



RhoA



PI3IK



Supplementary Figure 1.