

Sporadic Creutzfeldt-Jakob disease: Real-Time Quaking-Induced Conversion (RT-QuIC) assay represents a major diagnostic advance

Federico Angelo Cazzaniga*, Edoardo Bistaffa*, Chiara Maria Giulia De Luca, Antonio Indaco, Giuseppe Bufano, Giorgio Giaccone, Fabio Moda

Fondazione IRCCS Istituto Neurologico Carlo Besta, Division of Neurology 5–Neuropathology, Milan, Italy

*These authors equally contributed to this work

ABSTRACT

Sporadic Creutzfeldt-Jakob disease (sCJD) is a rare and fatal neurodegenerative disorder with an incidence of 1.5 to 2 cases per million population/year. The disease is caused by a proteinaceous infectious agent, named prion (or PrP^{Sc}), which arises from the conformational conversion of the cellular prion protein (PrP^C). Once formed, PrP^{Sc} interacts with the normally folded PrP^C coercing it to undergo similar structural rearrangement. The disease is highly heterogeneous from a clinical and neuropathological point of view. The origin of this variability lies in the aberrant structures acquired by PrP^{Sc}. At least six different sCJD phenotypes have been described and each of them is thought to be caused by a peculiar PrP^{Sc} strain. Definitive sCJD diagnosis requires brain analysis with the aim of identifying intracerebral accumulation of PrP^{Sc} which currently represents the only reliable biomarker of the disease. Clinical diagnosis of sCJD is very challenging and is based on the combination of several clinical, instrumental and laboratory tests representing surrogate disease biomarkers. Thanks to the advent of the ultrasensitive Real-Time Quaking-Induced Conversion (RT-QuIC) assay, PrP^{Sc} was found in several peripheral tissues of sCJD patients, sometimes even before the clinical onset of the disease. This discovery represents an important step forward for the clinical diagnosis of sCJD. In this manuscript, we present an overview of the current applications and future perspectives of RT-QuIC in the field of sCJD diagnosis.

Key words: Sporadic Creutzfeldt-Jakob disease; olfactory mucosa; cerebrospinal fluid; neurodegeneration; peripheral biomarkers; prion; seeding aggregation assays.

Correspondence: Giorgio Giaccone, Fondazione IRCCS Istituto Neurologico Carlo Besta, Division of Neurology 5 – Neuropathology, Via G. Celoria 11, 20133 Milan, Italy. E-mail: giorgio.giaccone@istituto-besta.it

Contributions: FAC, EB, manuscript drafting, tables and figures preparation; CMGD, AI, GB, contribution to manuscript drafting; FM, GG, manuscript concept and critical revision. All the authors have read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

Conflict of interest: The authors declare that they have no competing interests, and all authors confirm accuracy.

Molecular and neuropathological classification of sCJD subtypes

Among human prion diseases, sporadic Creutzfeldt-Jakob disease (sCJD) is the most common form affecting 1-2 individuals/million per year with similar distribution in males and females.¹ The age at onset is most frequently between 55 and 75 years.² sCJD presents with variable disease subtypes characterized by peculiar clinical and neuropathological features. In the past, other than the classical and more common subtypes, some clinical variants such as the Heidenhain, the myoclonic, the thalamic, the cerebellar or ataxic, and the panencephalopathic forms were reported.³⁻⁶ In general, sCJD cases present as multifocal and rapidly progressive encephalopathies with dementia, cerebellar ataxia, myoclonus while the progression of the disease results in an akinetic and mute state and the death occurs generally within 6 months after the disease onset.¹ The common mechanism underlying these pathologies is the spontaneous conformational conversion of the cellular prion protein (PrP^C) into an abnormally folded conformer named prion or PrP^{Sc}. This latter propagates in an autocatalytic manner in the brain by converting the PrP^C into the pathological isoform.

PrP^C is a glycosylphosphatidylinositol (GPI) anchored protein highly expressed in the central nervous system (CNS) and encoded by the *PRNP* gene located on chromosome 20 in humans.^{7,8} After its synthesis in the rough endoplasmic reticulum, PrP^C undergoes post-translational modifications comprising the C-terminal addition of the GPI anchor, the formation of a disulfide bridge between two C-terminal cysteine residues (Cys179-Cys214) and the N-linked glycosylation at asparagine residues (Asn181-Asn197).⁹

These oligosaccharides are further modified in the Golgi apparatus to produce complex-type chains enriched in sialic acid important for the synaptic localization of PrP^C.^{10,11} The different degrees of PrP^C glycosylation give rise to three isoforms of the protein: the di-glycosylated (70%), the mono-glycosylated (25%) and the un-glycosylated (5%) species.^{10,12} All these isoforms are rich in α -helices structures, soluble in detergent and are sensitive to proteolytic digestion with proteinase K (PK). Conversely, PrP^{Sc} is less soluble in detergent, has higher amount of β -sheet structures and is partially resistant to PK digestion. The limited proteolysis leads to the generation of N-terminal truncated fragments of di-, mono- and un-glycosylated PrP^{Sc} that migrate at lower molecular weights compared to those of PrP^C.¹³ Moreover, the un-glycosylated band of PrP^{Sc} can acquire two distinct molecular weights: 21 or 19 kDa which are referred to as type 1 or type 2 PrP^{Sc}, respectively.¹⁴ Neuropathologically, the main hallmarks of sCJD are spongiform changes, astrogliosis, neuronal loss and accumulation of PrP^{Sc} (Figure 1 and Figure 2).¹⁵

At present, PrP^{Sc} is the only disease-specific biomarker for sCJD and the definite diagnosis can be formulated post-mortem by biochemical and neuropathological analyses aimed at identifying the PrP^{Sc} accumulation in the CNS (Figure 2).^{16,17}

It is well known that PrP^{Sc} can acquire different abnormal conformations, named strains. The peculiar conformation of each strain can be faithfully transmitted to the host PrP^C and are believed to be responsible for the heterogeneity of prion diseases, in terms of tissue tropism, incubation period, clinical signs, neuropathological changes and interspecies transmission properties.¹⁸⁻²¹ In 1999, Parchi and colleagues²² classified sCJD in six major subtypes by correlating the clinical manifestations with the polymorphisms at codon 129 of the *PRNP* gene, *i.e.* methionine (M) or




Figure 1. Creutzfeldt-Jakob disease, hallmark neuropathologic lesions. Spongiform changes may appear as small vacuoles (A) diffusely present in grey matter (H&E, cerebral cortex, 20x) or large, confluent vacuolar lesions (B) typical of the MM2-C (cortical) subtype (H&E, cerebral cortex, 20x). Kuru plaques (C) small aggregates of PrP with the tinctorial and optical properties of amyloid are typically found in the cerebellum in MV2 subtype (H&E, 60x). Astrogliosis (D) may be severe in all subtypes of Creutzfeldt-Jakob disease (glial fibrillary acidic protein immunohistochemistry, 10x). Neuronal loss (E) is usually very severe in the cerebral cortex, basal ganglia and cerebellum (H&E, cerebellum) but may be mild in some cases (F) (microtubule associated protein 2 immunohistochemistry; 10x).

valine (V), and the electrophoretic mobility of the un-glycosylated PrP^{Sc} isoform in the brain after digestion with PK (type 1 or type 2 PrP^{Sc}). These findings demonstrated that the presence of M or V at codon 129 of PrP^C, as well as other still unknown factors, could modulate the structural rearrangement of PrP^C during misfolding, thus promoting the PrP^{Sc} strains variability.²³⁻²⁶ In addition, compelling evidence suggests that, in some sCJD cases, the CNS contains a mixture of PrP^{Sc} strains (e.g., MM1+2, VV1+2 and MV1+2), which make the classification of the disease even more challenging (as discussed in the next paragraphs).^{14,27-29}

The main pathological characteristics of each sCJD subtype are summarized in Table 1.

MM1 and MV1 subtype

MM1 is the most common form of sCJD (67% of all cases) while MV1 cases are rare (3%). Western blot analysis shows, for both subtypes, type 1 PrP^{Sc} and a glycoform pattern characterized by the predominance of the mono-glycosylated band. Despite the difference at codon 129 of *PRNP*, MM1 and MV1 cases share many pathological features. MM1/MV1 sCJD patients present with the myoclonic (or classic CJD) and the Heidenhain's variant.²² The mean age at onset of the disease is 66 years with an average clinical duration of 4 months. Clinical manifestations include cognitive impairment with memory loss and confusion/disorientation, depression, anxiety, psychosis and gait or limb ataxia.³⁰ Neuropathologically, the brain of these patients shows spongiosis with fine vacuoles. The basal ganglia, thalamus and cerebellum are less affected than the cerebral neocortex. The hippocampal cortex and brain stem are largely spared. The pattern of PrP^{Sc} deposition is synaptic and mainly affects the cerebral cortex while the cerebellum, the basal ganglia and thalamus are less involved (Figure 2 A,B).³¹ Moreover, the amount of PrP^{Sc} signal directly correlates with the severity of spongiosis.

VV2 subtype

The VV2 subtype corresponds to the cerebellar or ataxic variant and occurs in 15% of sCJD cases. The Western blot profile shows type 2 PrP^{Sc} with a preponderance of the mono-glycosylated isoform. The mean age at onset is 64 years (with a range of 40-83 years) and the clinical duration is about 7 months.³² Ataxia is the commonest early clinical feature accompanied by cognitive impairment and oculomotor signs while myoclonus is less frequent. In the late stages of the disease patients exhibit dementia, myoclonus and pyramidal signs. Neuropathologically, the spongiosis preferentially affects the deep layers of the frontal and occipital cortex, the entorhinal cortex and the hippocampus.³¹ Cerebral neocortex may be relatively spared particularly in cases with rapid courses. The cerebellar cortex is atrophic, with abundant PrP^{Sc} deposits characterized by a focal and plaque-like pattern that are negative for Congo Red and Thioflavin-S (amyloid stains). In addition, strong PrP^{Sc} deposition often occurs around neuronal perikarya in the cerebral cortex (Figure 2 C,D). The distribution of PrP^{Sc} immunostaining is affected by the disease duration. In cases with shorter disease duration, PrP^{Sc} involve diffusely the gray-matter region except for the neocortex which is affected only in patients with longer disease duration.³³

MV2 subtype

MV2 sCJD subtype is phenotypically and biochemically similar to VV2 cases (type 2 PrP^{Sc} and predominance of the mono-glycosylated form) and accounts for 10% of all sCJD. The mean age at onset is 65 years with a range of 36-83 years while the disease




Figure 2. Creutzfeldt-Jakob disease, patterns of pathological PrP deposition (PrP immunohistochemistry using 3F4 monoclonal antibody). In MM/MV1 subtype PrP deposition is mainly of synaptic type and appears as homogeneous, finely granular immunoreactivity in the neuropil of the cerebral cortex (A) (10x), while in the cerebellum is finely granular in the molecular layer while forming coarser deposits in the granular layer (B) (10x). In VV2 subtype PrP deposition often decorates the boundaries of pyramidal neurons of the cerebral cortex (C) (10x), while in the cerebellum plaque-like deposition takes is common in the granular layer (D) (10x). In most of the patients with the MM2 subtype PrP deposition takes variable aspects in the cerebral cortex (E) (20x), while the typical feature is the presence of Kuru plaques, small aggregates of PrP with the tinctorial and optical properties of amyloid, in the cerebellum (F) (10x). MM2-C (cortical) subtype is characterized by PrP immunoreactivity more intense at the rims of the large vacuoles of spongiosis in the cerebral cortex (G) (20x), while the cerebellum is usually relatively spared (H) (10x).

duration is significantly longer than VV2 with a mean of 17 months (range of 5-72 months). Early manifestations of the disease include oculomotor abnormalities, memory loss, behavioral disturbances and signs of the peripheral nervous system or medullary involvement. In MV2 cases ataxia is the most common presenting sign and cognitive deterioration, myoclonus and pyramidal signs, aphasia and apraxia arise during disease progression. The main hallmark that distinguishes MV2 cases from VV2 is the presence of cerebellar Kuru-type amyloid plaques in the Purkinje cell layer (positive for Congo Red and Thioflavin-S) (Figure 2 E,F).

MM2-thalamic subtype

MM2-thalamic (MM2T) sCJD is a rare subtype (2% of all the cases) known also as sporadic fatal insomnia (sFI).^{34,35} Biochemical characterization shows type 2 PrP^{Sc} with a predominant mono-glycosylated isoform. The mean age at onset is around 52 years (range 26-71 years) with a mean duration of 16 months (range 8-36 months). Besides insomnia, other common symptoms include dementia, and motor signs as ataxia, dysarthria, tremor, myoclonus and spasticity.^{34,36} Thalamus is the most affected brain region especially in the medial dorsal and anterior ventral nuclei where marked atrophy (observed also in the inferior olivary nucleus) and severe astrogliosis is accompanied by prominent neuronal loss. Spongiform changes and faint PrP^{Sc} deposition may be present in the cerebral cortex.

MM2-cortical subtype

This rare cortical variant of MM2 subtype (type 2 PrP^{Sc} with a prevalence of the mono-glycosylated band) represents 2% of all sCJD cases and is characterized by progressive dementia and disturbances of higher cognitive functions, high-frequency aphasia and apraxia and late myoclonus or epileptic seizures.³⁷ The average age at onset is 64 years with a 49-77 year range and the disease duration is approximately 16 months. Brain lesions are similar to

that of MM1 or MV1 subtypes but, despite the relatively long disease duration, the cerebellum is almost spared. Large vacuoles are present in the cerebral cortex, basal ganglia and thalamus which might be confluent. Immunodetection of PrP^{Sc} reveals a coarse pattern of staining which occasionally localizes at the rim of the vacuoles (perivacuolar PrP^{Sc} deposition) (Figure 2 G,H).³¹

VV1 subtype

VV1 is the rarest subtype of sCJD representing 1% of the total cases. The Western blot analysis shows type 1 PrP^{Sc} with a prevalence of the mono-glycosylated isoform. Patients are relatively younger (mean age at onset 44 years) compared to other sCJD subtypes with a mean duration of 21 months (range 17-42 months). Early symptoms include psychiatric or cognitive abnormalities that evolve in extrapyramidal signs and ataxia while myoclonus was observed only in few patients. Massive spongiform lesions affect the cortico-striatal regions while other subcortical regions and cerebellum are almost spared. Although the severe spongiform changes observed in VV1 patients, PrP^{Sc} immunohistochemistry shows faint punctate staining confined in the cerebral cortex.³¹

Mixed subtypes

Type 1 and type 2 PrP^{Sc} have been found to co-exist in about 35% of sCJD cases and may be present in the same or distinct anatomical regions of the same patient.²⁹ This finding is more frequent in MM (43%) than MV (23%) and VV (15%) cases.¹⁴ The predominance of PrP^{Sc} type 1 or 2 influences the clinical and neuropathological phenotype of the diseases. The MM1+2 cases mimic the clinical phenotype of MM1 while the PrP^{Sc} deposition is a combination of the typical neuropathological features of MM1 and MM2 (synaptic and perivacuolar patterns, respectively). Conversely, VV1+2 subjects are similar to VV2 sCJD cases in terms of clinical and neuropathological features.¹⁴

Table 1. Pathological features of sCJD molecular subtypes.

sCJD molecular subtypes	% of cases	Median age at onset (years)	Duration (months)	Main neuropathological alterations
MM1	67	66	~ 4	Diffuse spongiosis with small vacuoles affecting the neocortex, striatum and cerebellar cortex. Synaptic pattern of PrP ^{Sc} deposition
MV1	3	66	~ 4	
VV1	1	44	~ 21	Severe spongiosis with fine vacuoles in the cerebral cortex and striatum. Punctate pattern of PrP ^{Sc} deposition
MM2 - thalamic	2	52	~16	Atrophy of the thalamus and inferior olivary nuclei with spongiform alterations confined to the cerebral cortex. Weak and synaptic pattern of PrP ^{Sc} deposition.
MM2 - cortical	2	64	~ 16	Severe spongiosis with large confluent vacuoles predominantly in cerebral cortex and striatum. Perivacuolar and coarse pattern of PrP ^{Sc} deposition.
MV2	10	65	~ 17	Diffuse and confluent spongiosis similar to VV2 subtype. Amyloid Kuru plaques in the molecular and granular layer of the cerebellum.
VV2	15	64	~ 7	Spongiform changes found in the cerebellum, striatum, thalamus and brainstem. Plaque-like and perineuronal pattern of PrP ^{Sc} deposition.

MM1, Methionine/Methionine – PrP^{Sc} type 1; MV1, Methionine/Valine - PrP^{Sc} type 1; VV1, Valine/Valine - PrP^{Sc} type 1; MM2-T, Methionine/Methionine – Thalamic PrP^{Sc} type 2; MV2-C, Methionine/Valine – Cortical - PrP^{Sc} type 2; MV2, Methionine/Valine PrP^{Sc} type 2; VV2, Valine/Valine PrP^{Sc} type 2.

Clinical challenges

The clinical diagnosis of sCJD is particularly challenging especially in the early stages of the disease.³⁸ It relies on defined criteria that classify the disease as possible or probable.³⁹ Several clinical, instrumental and laboratory tests are commonly used to formulate an *in vivo* diagnosis of sCJD: electroencephalogram (EEG),⁴⁰ magnetic resonance imaging (MRI)⁴¹ and cerebrospinal fluid (CSF) biomarkers analysis. Several CSF biomarkers have been investigated including the 14-3-3 protein, total tau (t-tau) and phosphorylated tau (p-tau) proteins, neurofilament light chain (NfL), neuron-specific enolase and α -synuclein. The most reliable and commonly used are 14-3-3 and t-tau.^{42,43}

14-3-3 protein is a biomarker of neuronal cell death and therefore it is not specific for prion diseases. It is commonly reported to possess an average sensitivity of 85-95% and specificity of 40-100%^{32,44-48} for sCJD. However, the main issue in using the 14-3-3 as a biomarker for prion diseases lies in the fact that its elevation is common in some neurologic and neurodegenerative diseases including herpes simplex encephalitis, other encephalitis, intracerebral metastases, metabolic encephalopathy, hypoxic brain damage, dementia with Lewy bodies (DLB) and Alzheimer's disease (AD).^{33,49-51} Therefore, 14-3-3 analysis may increase the probability of CJD when other clinical features are suggestive of prion disease but it cannot be assumed as a specific biomarker.⁵²

Increased levels of t-tau (cut off >1300 pg/mL) may identify sCJD patients with a sensitivity of 67-91% and a specificity of 67-95%.^{44,46,47,53-56} This measurement helps to differentiate sCJD from AD. Indeed, t-tau was 3.1 times higher in sCJD compared to AD and 41 times higher than in healthy subjects.⁵⁷ Recently, the ratio t-tau/p-tau was found elevated in sCJD patients with a specificity of 94-97% and a sensitivity ranging from 75-94%.^{46,58-60}

Among other CSF biomarkers proposed for prion disease diagnosis, NfL has been reported to be significantly elevated in sCJD compared to other neurodegenerative disorders like AD, DLB, frontotemporal dementia and vascular dementia. However, despite increased NfL levels enable discrimination of sCJD from normal controls,⁶¹⁻⁶⁴ they do not consent accurate discrimination between sCJD and other rapidly progressive dementias,⁶⁵ neurodegenerative dementia⁶⁴ and neurological diseases with dementia syndromes.⁶² Recently, serum NfL analysis has been suggested as a diagnostic marker for prion diseases showing similar sensitivity and specificity to CSF markers in differentiating sCJD from healthy subjects.^{66,67}

α -synuclein (α -syn) is commonly used as a biomarker for a group of diseases known as α -synucleinopathies, which includes, among the others, Parkinson's disease (PD)⁶⁸ and dementia with Lewy bodies (DLB),⁶⁹ but its usefulness for CJD diagnosis has been recently investigated. Two studies reported that total α -syn (t- α -syn) was specifically elevated in CSF of sCJD patients compared to control subjects.^{70,71} Similarly, the phospho-serine-129 α -synuclein (p- α -syn) was found elevated in the CSF of sCJD patients compared to PD, DLB and neurological controls. A combined analysis of both markers, showed 90.5% sensitivity and 97.6% specificity for sCJD diagnosis.⁷² Other CSF and serum biomarkers of prion diseases, including the neuron specific enolase (NSE),^{73,74} the S100B protein,⁷⁵ SERPINA3⁷⁶ and thymosin β 4⁷⁷ are currently under investigation. Unfortunately, although useful for the clinical diagnosis of CJD, CSF biomarkers are not disease-specific.

The definite diagnosis depends on *post-mortem* examination of the brain aimed at identifying and characterizing the disease-specific biomarker of prion diseases, the PrP^{Sc}. Through a combination of biochemical (*e.g.*, Western blot after PK digestion), immunohistochemical and genetic analyses it is possible to identify the specific sCJD subtype. Thanks to the recent development of

the ultrasensitive seeding aggregation assays, named Real-Time Quaking Induced Conversion (RT-QuIC) and Protein Misfolding Cyclic Amplification (PMCA) the diagnostic accuracy of prion diseases has been significantly increased. In particular, the PMCA enabled efficient detection of traces of PrP^{Sc} in the CSF, urine and blood of patients with variant CJD (vCJD), which is related to the consumption of foodstuff obtained from cattle affected by bovine spongiform encephalopathy. However, this technique, has never been able to efficiently detect PrP^{Sc} associated with sCJD.⁷⁸⁻⁸¹ In contrast, the RT-QuIC has been optimized to efficiently detect low amounts of sCJD prions in the CSF, olfactory mucosa and skin samples in a more rapid and safe manner (with respect to PMCA) while requiring a limited handling of the specimens and reducing the risk of their contamination.⁸²⁻⁸⁵ For this reason, the RT-QuIC has been adopted by several specialized centers for the analysis of biological samples collected from patients with suspected sCJD, as detailed in the next section.




Figure 3. Schematic illustration of the RT-QuIC reaction. The RT-QuIC process is divided into three phases: (1) the lag phase, (2) the growth phase and (3) the plateau phase. The reaction mix is composed by recombinant PrP (recPrP) and Thioflavin T (ThT) which are dissolved in common buffers. The addition of PrP^{Sc} (pink triangle) to the reaction induces the conversion of recPrP (blue hexagon) into a misfolded form (red arrow) which starts to aggregate and form recPrP amyloid fibrils. In the absence of PrP^{Sc}, recPrP can aggregate (dotted line) following a well-defined kinetics. The formation of the aggregates induces the emission of a ThT fluorescence signal (yellow star). In the presence of PrP^{Sc}, the kinetics of recPrP aggregation is significantly accelerated (solid line). The increased kinetics of recPrP aggregation is known as seeding effect.

RT-QuIC assay

RT-QuIC is an ultrasensitive technique developed by Atarashi *et al.* in 2011 in the field of prion diseases. This assay exploits the intrinsic ability of PrP^{Sc} to promote the conformational rearrangement of PrP^C that can aggregate into amyloid fibrils.⁸⁶ The assay mimics *in vitro* the process of PrP^C misfolding and aggregation which occurs *in vivo*. Recombinant PrP^C (recPrP) with the amino acid sequence of different species can be used as a reaction substrate. The addition of traces of PrP^{Sc} to the reaction substrate induces its aggregation and the kinetics of this process can be monitored in real-time by using a fluorescent dye, named Thioflavin-T (ThT).⁸⁷ In general, each sample is analyzed in quadruplicates using a multi-well plate.⁸⁸ The samples are subjected to cyclic phases of incubation and shaking using a dedicated fluorescence microplate reader.⁸⁹ In the presence of PrP^{Sc}, the incubation phase stimulates the formation of recPrP amyloid fibrils, while the shaking phase permits the fragmentation of the aggregates into smaller units capable to recruit and convert further recPrP into new amyloid fibrils.⁹⁰

The *in vitro* aggregation process can be represented on a cartesian plane where fluorescence is plotted against time generating a kinetic curve characterized by three phases: i) a lag phase, where PrP^{Sc} interacts with recPrP and induces this latter to misfold ii) a growth phase, where misfolded recPrP aggregate to form oligomers and small amyloid fibrils sensitive to ThT (exponential

increase of fluorescence) and iii) a plateau phase, where almost all recPrP is incorporated into fibrils. Under normal reaction conditions, recPrP spontaneously aggregates while the addition of PrP^{Sc} (even in traces) to the substrate significantly accelerates the kinetic of recPrP aggregation (seeding effect) (Figure 3). A sample is considered positive when at least 2 out of 4 replicates show a seeding effect. The RT-QuIC end-products are partially resistant to PK digestion.⁹¹

RT-QuIC enabled PrP^{Sc} detection in CSF, olfactory mucosa (OM), skin, eye, peripheral nerve, and digestive system of patients with different forms of prion diseases (Table 2).

The assay developed in 2011 was considered the “first generation RT-QuIC” since the analyses were performed at 42°C using the recombinant full-length Syrian Hamster prion protein (recSHA₍₂₃₋₂₃₁₎) as reaction substrate. With this experimental setting it was possible to detect PrP^{Sc} in the CSF of a series of Japanese subjects with sCJD and 30 Australian sCJD patients with 80% sensitivity and 100% specificity.⁹²

One year later, the analyses of 123 patients with neuropathologically confirmed sCJD showed that RT-QuIC was able to identify PrP^{Sc} in CSF with a sensitivity of 91% and specificity of 98%.⁸⁴

In 2014, Orrù and colleagues⁸⁴ performed RT-QuIC analysis of OM and CSF collected from living patients with possible or probable clinical diagnosis of CJD. The RT-QuIC analysis of OM identified 30 out of 31 sCJD patients with a sensitivity of 97% and specificity of 100% while the analysis of CSF showed less sensi-

Table 2. Specificity and sensitivity of 1st and 2nd generation of RT-QuIC.

Samples	Year	Reference	Substrate recPrP	Sensitivity %	Specificity %
CSF	2011	Atarashi <i>et al.</i> ⁸²	recSHA (23-231)	91.5	100.0
	2012	McGuire <i>et al.</i> ⁸⁸	recSHA (23-231)	89.0	99.0
	2014	Orrù <i>et al.</i> ⁸⁴	recSHA (23-231)	77.0	100.0
	2015	Cramm <i>et al.</i> ¹¹⁴	recSHA (23-231)	Not reported	Not reported
	2015	Orrù <i>et al.</i> ¹¹⁵	recSHA (90-231)	95.8	100.0
	2016	Cramm <i>et al.</i> ⁹³	recSHA (23-231)	85.0	99.0
	2016	Groveman <i>et al.</i> ⁹⁴	recSHA (23-231)	72.5	100.0
	2016	Groveman <i>et al.</i> ⁹⁴	recSHA (90-231)	93.8	100.0
	2016	Park <i>et al.</i> ⁹⁵	recSHA (23-231)	76.5	100.0
	2016	McGuire <i>et al.</i> ⁹²	recSHA (23-231)	100.0	100.0
	2017	Franceschini <i>et al.</i> ⁸³	recSHA (90-231)	97.2	100.0
	2017	Bongianni <i>et al.</i> ⁹⁶	recSHA (23-231)	71.4	100.0
	2017	Bongianni <i>et al.</i> ⁹⁶	recSHA (90-231)	82.6	100.0
	2017	Lattanzio <i>et al.</i> ⁵⁶	recSHA (23-231)	82.1	99.4
	2017	Foutz <i>et al.</i> ¹⁰⁷	recSHA (90-231)	92.0	98.5
	2017	Foutz <i>et al.</i> ¹⁰⁷	recSHA (90-231)	95.0	100.0
	2018	Rudge <i>et al.</i> ¹¹⁰	recSHA (23-231)	89.0	100.0
	2018	Hermann <i>et al.</i> ¹⁰⁹	recSHA (23-231)	97.0	99.0
	2019	Abu-Rumeileh <i>et al.</i> ⁵⁵	recSHA (23-231)	82.5	100.0
	2019	Abu-Rumeileh <i>et al.</i> ⁵⁵	recSHA (90-231)	97.4	100.0
2020	Fiorini <i>et al.</i> ¹⁰⁰	recSHA (90-231)	96.0	100.0	
2020	Rhoads <i>et al.</i> ⁹⁷	recSHA (90-231)	90.3	98.5	
2020	Xiao <i>et al.</i> ⁹⁸	recSHA (90-231)	96.7	100.0	
OM	2014	Orrù <i>et al.</i> ⁸⁴	recSHA (23-231)	97.0	100.0
	2017	Bongianni <i>et al.</i> ⁹⁶	recSHA (90-231)	92.0	100.0
	2020	Fiorini <i>et al.</i> ¹⁰⁰	recSHA (90-231)	91.4	100.0
Skin	2017	Orrù <i>et al.</i> ¹⁰⁵	recSHA (23-231)	100.0	100.0
	2020	Mammana <i>et al.</i> ⁸⁵	recSHA (23-231)	89.0	100.0
Eye	2018	Orrù <i>et al.</i> ¹⁰⁶	recSHA (90-231)	100.0	100.0
PN	2019	Baiardi <i>et al.</i> ¹¹⁶	recSHA (90-231)	100.0%	100.0%
DS	2019	Satoh <i>et al.</i> ¹¹⁷	not reported	100.0%	not reported

CSF, cerebrospinal fluid; OM, olfactory mucosa; PN, peripheral nerve; DS, digestive system; recSHA₍₂₃₋₂₃₁₎, recombinant full-length syrian hamster prion protein; recSHA₍₉₀₋₂₃₁₎, recombinant N-terminally truncated syrian hamster prion protein.

tivity (77%) but similar specificity (100%). In 2016, the first multi-center studies demonstrated the reproducibility, reliability, and robustness of the first generation of CSF RT-QuIC (PQ-CSF) in clinical practice.^{92,93}

In 2015, the group of Caughey optimized the RT-QuIC protocol and developed the “second generation RT-QuIC” by introducing two modifications: i) the use of N-terminally truncated recSHA_(90–231) as a reaction substrate and ii) the increasing of temperature from 42°C to 55°C. The use of a different substrate and a higher temperature improved the RT-QuIC performances by identifying positive CSF samples in shorter time (4–14 hours) compared to those required by the first generation assay (2,5–5 days). In particular, the RT-QuIC analysis identified PrP^{Sc} in 46 out of 48 CSF samples collected from sCJD individuals yielding 95.8% diagnostic sensitivity and 100% specificity. Another retrospective study of Groveman *et al.*⁹⁴ confirmed the high sensitivity and specificity (94% and 100%, respectively) of the CSF RT-QuIC test in a large cohort of patients (n=113) with probable or definitive sCJD diagnosis. Further investigations were then performed with the aim of improving the sensitivity of the assay in detecting PrP^{Sc} in CSF and OM.^{55,56,83,92,95–98} In 2020, an international trial confirmed the robustness and reliability of the second generation RT-QuIC for the diagnosis of sCJD.⁹⁹ Furthermore, a recent article by Fiorini *et al.*¹⁰⁰ demonstrated that through the combined RT-QuIC analysis of the CSF and OM collected from the same sCJD patient it is possible to reach a diagnostic accuracy of 100%. For these reasons, the RT-QuIC has been introduced among the diagnostic criteria of some surveillance centers.¹⁰¹ Recent evidence showed that recPrP with the bank vole amino acid sequence can detect almost all PrP strains (from human or animal origin), but it has not been introduced in the diagnostic field yet.^{102–104}

From 2017, the RT-QuIC assay has been extended to the analysis of other peripheral tissues. In particular, Orrù *et al.* explored the potential prion seeding activity and infectivity of skin collected from 21 sCJD patients and eye (retina, sclera and cornea) collected *post-mortem* from 11 sCJD cases.^{105,106} Similarly, Mammama *et al.* analyzed skin biopsies of sCJD patients and identified PrP^{Sc} with 89% sensitivity and 100% specificity.⁸⁵

Correlations of RT-QuIC results with neuropathological findings

To date, only few studies have investigated whether there is a correlation between the RT-QuIC results and the phenotypes of sCJD.^{107,108} In the case of CSF samples, the sensitivity of RT-QuIC was found to be high in the most common MM1/MV1 and VV2 sCJD cases, while it was lower in MV2 cases (75–93%).^{55,56,83,97,107,109} In other rare subtypes, including VV1 and MM2, the sensitivity was found to range between 0–100% and 44–78%, respectively.^{56,97,109–111} In these latter cases, the limited amount of CSF, hampered the possibility to properly evaluate the diagnostic accuracy of the assay. In 2016, Foutz *et al.* observed a correlation between RT-QuIC kinetics and sCJD subtypes. In particular, they observed that MM1 cases had significantly shorter lag phase and higher fluorescence values compared to MM2 cases, and these findings enabled discrimination of both phenotypes with an accuracy of 95%. At the same time, the extended lag phase and lower intensity of fluorescence allowed to differentiate VV1 to VV2 individuals with an accuracy of 80%. MV1, MV2, and mixed type cases did not show significant differences in terms of lag phases or fluorescence intensities.¹⁰⁷ Recently, Piconi *et al.* subjected to PK digestion the RT-QuIC products obtained from the analysis of brain homogenates (BH) and CSF of patients with the six phenotypes of sCJD. In this case, regardless of the sCJD subtype, all

samples displayed PK-resistant signal characterized by similar electrophoretic mobility and banding profile, even when challenged with several anti-PrP antibodies.¹⁰⁸ Thus, in contrast to the work of Foutz, they could not identify peculiar features useful to distinguish the six sCJD subtypes. For this reason, the possibility to identify sCJD subtypes by RT-QuIC remains to be clearly elucidated. Very recent findings show that formalin fixed brains are capable to exert an efficient seeding activity by RT-QuIC, using both animal¹¹² and human specimens (*personal communication*).

Conclusions

Currently, the RT-QuIC test represents the most reliable and powerful tool for the early detection of PrP^{Sc} in peripheral tissues of patients with a suspected clinical diagnosis of sCJD.⁹⁰ The reason for the rapid growth of RT-QuIC use in the clinical practice, although still confined to specialized laboratories, lays in the fact that it is not invasive for the patients, has a relatively low cost and a high predictive value. Among the advantages, the method is not time-consuming and enables the analysis of a huge number of samples in a relatively short period of time.¹¹³ Overall, these characteristics support the choice by WHO to include the CSF RT-QuIC test in the diagnostic criteria for sCJD.^{39,99,101} As previously mentioned, only few specialized laboratories have adopted the RT-QuIC technology. However, the assay is relatively easy to learn and can be rapidly used by trained personnel, thus consenting its widening in other centers specialized in the diagnosis of neurodegenerative diseases associated with protein misfolding. Future multi-center trials will consent to verify the robustness of the RT-QuIC for the analysis of new peripheral tissues (*e.g.* OM, skin) and to further explore the potential of this assay to stratify patients in their early disease stage.

References

- Ladogana A, Puopolo M, Croes EA, Budka H, Jarius C, Collins S, et al. Mortality from Creutzfeldt-Jakob disease and related disorders in Europe, Australia, and Canada. *Neurology* 2005;64:1586–91.
- Imran M, Mahmood S. An overview of human prion diseases. *Virology* 2011;8:559.
- Heidenhain A. [Klinische und anatomische Untersuchungen über eine eigenartige organische Erkrankung des Zentralnervensystems im Praesenium]. [Article in German]. *Z Gesamte Neurol Psy* 1929;118:49–114.
- Mittal M, Hammond N, Husmann K, Lele A, Pasnoor M. Creutzfeldt-Jakob disease presenting as bulbar palsy. *Muscle Nerve* 2010;42:833–5.
- Alema G, Bignami A. [Subacute degenerative presenile polioencephalopathy with akinetic stupor and decorticate rigidity with myoclonus (“myoclonic” variety of the Jakob-Creutzfeld disease)]. [Article in Italian]. *Riv Sper Freniatr Med Leg Alien Ment* 1959;83:S1485–623.
- Nowacki P, Kulczycki J, Narolewska A, Grzelec H. Amyotrophic form of Creutzfeldt-Jakob disease with rapid course in 82-year-old man. *Folia Neuropathol* 2000;38:161–3.
- Stahl N. Scrapie prion protein contains a phosphatidylinositol glycolipid. *Cell* 1987;51:229–40.
- Castle AR, Gill AC. Physiological functions of the cellular prion protein. *Front Mol Biosci* 2017;4:19.
- Turk E, Teplow DB, Hood LE, Prusiner SB. Purification and properties of the cellular and scrapie hamster prion proteins.

- Eur J Biochem 1988;176:21–30.
10. Caughey B, Race RE, Ernst D, Buchmeier MJ, Chesebro B. Prion protein biosynthesis in scrapie-infected and uninfected neuroblastoma cells. *J Virol* 1989;63:175–81.
 11. Bate C, Nolan W, McHale-Owen H, Williams A. Sialic acid within the glycosylphosphatidylinositol anchor targets the cellular prion protein to synapses. *J Biol Chem* 2016;291:17093–101.
 12. Rudd PM, Endo T, Colominas C, Groth D, Wheeler SF, Harvey DJ, et al. Glycosylation differences between the normal and pathogenic prion protein isoforms. *Proc Natl Acad Sci USA* 1999;96:13044–9.
 13. Oesch B, Westaway D, Wälchli M, McKinley MP, Kent SBH, Aebersold R, et al. A cellular gene encodes scrapie PrP 27-30 protein. *Cell* 1985;40:735–46.
 14. Parchi P, Strammiello R, Notari S, Giese A, Langeveld JPM, Ladogana A, et al. Incidence and spectrum of sporadic Creutzfeldt–Jakob disease variants with mixed phenotype and co-occurrence of PrPSc types: an updated classification. *Acta Neuropathol* 2009;118:659–71.
 15. Soto C, Satani N. The intricate mechanisms of neurodegeneration in prion diseases. *Trends Mol Med* 2011;17:14–24.
 16. Prusiner SB. Prions. *Proc Natl Acad Sci USA* 1998;95:13363–83.
 17. Soto C. Diagnosing prion diseases: needs, challenges and hopes. *Nat Rev Microbiol* 2004;2:809–19. A
 18. Morales R, Hu PP, Duran-Aniotz C, Moda F, Diaz-Espinoza R, Chen B, et al. Strain-dependent profile of misfolded prion protein aggregates. *Sci Rep* 2016;6:20526.
 19. Rossi M, Baiardi S, Parchi P. Understanding prion strains: Evidence from studies of the disease forms affecting humans. *Viruses* 2019;11:309.
 20. Morales R, Abid K, Soto C. The prion strain phenomenon: Molecular basis and unprecedented features. *Biochim Biophys Acta* 2007;1772:681–91.
 21. Zafar S, Younas N, Shafiq M, Zerr I. Prion protein strain diversity and disease pathology. in: prions - Some physiological and pathophysiological aspects IntechOpen; 2019.
 22. Parchi P, Giese A, Capellari S, Brown P, Schulz-Schaeffer W, Windl O, et al. Classification of sporadic Creutzfeldt–Jakob disease based on molecular and phenotypic analysis of 300 subjects. *Ann Neurol* 1999;46:224–33.
 23. Hosszu LLP, Jackson GS, Trevitt CR, Jones S, Batchelor M, Bhelt D, et al. The residue 129 polymorphism in human prion protein does not confer susceptibility to Creutzfeldt–Jakob disease by altering the structure or global stability of PrPc. *J Biol Chem* 2004; 279:28515–21.
 24. Safar JG. Molecular pathogenesis of sporadic prion diseases in man. *Prion* 2012;6:108–15.
 25. Kobayashi A, Iwasaki Y, Otsuka H, Yamada M, Yoshida M, Matsuura Y, et al. Deciphering the pathogenesis of sporadic Creutzfeldt–Jakob disease with codon 129 M/V and type 2 abnormal prion protein. *Acta Neuropathol Commun* 2013;1:74.
 26. Kobayashi A, Teruya K, Matsuura Y, Shirai T, Nakamura Y, Yamada M, et al. The influence of PRNP polymorphisms on human prion disease susceptibility: an update. *Acta Neuropathol* 2015;130:159–70.
 27. Uro-Coste E, Cassard H, Simon S, Lugan S, Bilheude JM, Perret-Liaudet A, et al. Beyond PrPres type 1/type 2 dichotomy in Creutzfeldt–Jakob disease. *PLoS Pathog* 2008;4:e1000029.
 28. Cassard H, Huor A, Espinosa JC, Douet JY, Lugan S, Aron N, et al. Prions from sporadic Creutzfeldt–Jakob disease patients propagate as strain mixtures. *MBio* 2020;11:e00393–20.
 29. Puoti G, Giaccone G, Rossi G, Canciani B, Bugiani O, Tagliavini F. Sporadic Creutzfeldt–Jakob disease: Co-occurrence of different types of PrPSc in the same brain. *Neurology* 1999;53:2173–6.
 30. Zerr I, Schulz-Schaeffer WJ, Giese A, Bodemer M, Schröter A, Henkel K, et al. Current clinical diagnosis in Creutzfeldt–Jakob disease: Identification of uncommon variants. *Ann Neurol* 2000;48:323–9.
 31. Parchi P, de Boni L, Saverioni D, Cohen ML, Ferrer I, Gambetti P, et al. Consensus classification of human prion disease histotypes allows reliable identification of molecular subtypes: an inter-rater study among surveillance centres in Europe and USA. *Acta Neuropathol* 2012;124:517–29.
 32. Collins SJ. Determinants of diagnostic investigation sensitivities across the clinical spectrum of sporadic Creutzfeldt–Jakob disease. *Brain* 2006;129:2278–87.
 33. Baiardi S, Magherini A, Capellari S, Redaelli V, Ladogana A, Rossi M, et al. Towards an early clinical diagnosis of sporadic CJD VV2 (ataxic type). *J Neurol Neurosurg Psychiatry* 2017;88:764–72.
 34. Parchi P, Capellari S, Chin S, Schwarz HB, Schecter NP, Butts JD, et al. A subtype of sporadic prion disease mimicking fatal familial insomnia. *Neurology* 1999;52:1757–63.
 35. Puoti G, Bizzi A, Forloni G, Safar JG, Tagliavini F, Gambetti P. Sporadic human prion diseases: molecular insights and diagnosis. *Lancet Neurol* 2012;11:618–28.
 36. Scaravilli F, Cordery RJ, Kretschmar H, Gambetti P, Brink B, Fritz V, et al. Sporadic fatal insomnia: A case study. *Ann Neurol* 2000;48:665–9.
 37. Krasnianski A, Meissner B, Schulz-Schaeffer W, Kallenberg K, Bartl M, Heinemann U, et al. Clinical features and diagnosis of the MM2 Cortical subtype of sporadic Creutzfeldt–Jakob disease. *Arch Neurol* 2006;63:876.
 38. Mead S, Rudge P. CJD mimics and chameleons. *Pract Neurol* 2017;17:113–21.
 39. Zerr I, Kallenberg K, Summers DM, Romero C, Taratuto A, Heinemann U, et al. Updated clinical diagnostic criteria for sporadic Creutzfeldt–Jakob disease. *Brain* 2009;132:2659–68.
 40. Zerr I, Pocchiari M, Collins S, Brandel JP, de Pedro Cuesta J, Knight RSG, et al. Analysis of EEG and CSF 14-3-3 proteins as aids to the diagnosis of Creutzfeldt–Jakob disease. *Neurology* 2000;55:811–5.
 41. Collie DA, Sellar RJ, Zeidler M, Colchester ACF, Knight R, Will RG. MRI of Creutzfeldt–Jakob disease: Imaging features and recommended MRI protocol. *Clin Radiol* 2001;56:726–39.
 42. Karch A, Zerr I. A comparison of tau and 14-3-3 protein in the diagnosis of Creutzfeldt–Jakob disease. *Neurology* 2013;80:2081.
 43. Blennow K, Johansson A, Zetterberg H. Diagnostic value of 14-3-3 β immunoblot and T-tau/P-tau ratio in clinically suspected Creutzfeldt–Jakob disease. *Int J Mol Med* 2005;16:1147–9.
 44. Coulthart MB, Jansen GH, Olsen E, Godal DL, Connolly T, Choi BCK, et al. Diagnostic accuracy of cerebrospinal fluid protein markers for sporadic Creutzfeldt–Jakob disease in Canada: a 6-year prospective study. *BMC Neurol* 2011;11:133.
 45. Chohan G, Pennington C, Mackenzie JM, Andrews M, Everington D, Will RG, et al. The role of cerebrospinal fluid 14-3-3 and other proteins in the diagnosis of sporadic Creutzfeldt–Jakob disease in the UK: a 10-year review. *J Neurol Neurosurg Psychiatry* 2010;81:1243–8.
 46. Bahl JMC, Heegaard NHH, Falkenhorst G, Laursen H, Høgenhaven H, Mølbak K, et al. The diagnostic efficiency of biomarkers in sporadic Creutzfeldt–Jakob disease compared to

- Alzheimer's disease. *Neurobiol Aging* 2009;30:1834–41.
47. Sanchez-Juan P, Green A, Ladogana A, Cuadrado-Corrales N, Saanchez-Valle R, Mitrova E, et al. CSF tests in the differential diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2006;67:637–43.
 48. Stoeck K, Sanchez-Juan P, Gawinecka J, Green A, Ladogana A, Pocchiari M, et al. Cerebrospinal fluid biomarker supported diagnosis of Creutzfeldt-Jakob disease and rapid dementias: a longitudinal multicentre study over 10 years. *Brain* 2012;135:3051–61.
 49. Satoh J, Kurohara K, Yukitake M, Kuroda Y. The 14-3-3 protein detectable in the cerebrospinal fluid of patients with prion-unrelated neurological diseases is expressed constitutively in neurons and glial cells in culture. *Eur Neurol* 1999;41:216–25.
 50. Zerr I, Bodemer M, Gefeller O, Otto M, Poser S, Wiltfang J, et al. Detection of 14-3-3 protein in the cerebrospinal fluid supports the diagnosis of Creutzfeldt-Jakob disease. *Ann Neurol* 1998;43:32–40.
 51. Chapman T, McKeel DW, Morris JC. Misleading results with the 14-3-3 assay for the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2000;55:1396–8.
 52. Muayqil T, Gronseth G, Camicioli R. Evidence-based guideline: Diagnostic accuracy of CSF 14-3-3 protein in sporadic Creutzfeldt-Jakob disease: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology* 2012;79:1499–506.
 53. Hamlin C, Puoti G, Berri S, Sting E, Harris C, Cohen M, et al. A comparison of tau and 14-3-3 protein in the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2012;79:547–52.
 54. Former SA, Takada LT, Bettecher BM, Lobach I V, Tartaglia MC, Torres-Chae C, et al. Comparing CSF biomarkers and brain MRI in the diagnosis of sporadic Creutzfeldt-Jakob disease. *Neurol Clin Pract* 2015;5:116–25.
 55. Abu-Rumeileh S, Baiardi S, Polisch B, Mammanna A, Franceschini A, Green A, et al. Diagnostic value of surrogate CSF biomarkers for Creutzfeldt-Jakob disease in the era of RT-QuIC. *J Neurol* 2019;266:3136–43.
 56. Lattanzio F, Abu-Rumeileh S, Franceschini A, Kai H, Amore G, Poggiolini I, et al. Prion-specific and surrogate CSF biomarkers in Creutzfeldt-Jakob disease: diagnostic accuracy in relation to molecular subtypes and analysis of neuropathological correlates of p-tau and A β 42 levels. *Acta Neuropathol* 2017;133:559–78.
 57. Blennow K, Diaz-Lucena D, Zetterberg H, Villar-Pique A, Karch A, Vidal E, et al. CSF neurogranin as a neuronal damage marker in CJD: a comparative study with AD. *J Neurol Neurosurg Psychiatry* 2019;90:846–53.
 58. Skillbäck T, Rosén C, Asztely F, Mattsson N, Blennow K, Zetterberg H. Diagnostic Performance of cerebrospinal fluid total tau and phosphorylated tau in Creutzfeldt-Jakob disease. *JAMA Neurol* 2014;71:476.
 59. Baldeiras IE, Ribeiro MH, Pacheco P, Machado Á, Santana I, Cunha L, et al. Diagnostic value of CSF protein profile in a Portuguese population of sCJD patients. *J Neurol* 2009;256:1540–50.
 60. Skinningsrud A, Stenset V, Gundersen AS, Fladby T. Cerebrospinal fluid markers in Creutzfeldt-Jakob disease. *Cerebrospinal Fluid Res* 2008;5:14.
 61. Kovacs GG, Andreasson U, Liman V, Regelsberger G, Lutz MI, Danics K, et al. Plasma and cerebrospinal fluid tau and neurofilament concentrations in rapidly progressive neurological syndromes: a neuropathology-based cohort. *Eur J Neurol* 2017;24:1326–e77.
 62. Zerr I, Schmitz M, Karch A, Villar-Piqué A, Kanata E, Golanska E, et al. Cerebrospinal fluid neurofilament light levels in neurodegenerative dementia: Evaluation of diagnostic accuracy in the differential diagnosis of prion diseases. *Alzheimer's Dement* 2018;14:751–63.
 63. Antonell A, Tort Merino A, Ríos J, Balasa M, Borrego Écija S, Auge JM, et al. Synaptic, axonal damage and inflammatory cerebrospinal fluid biomarkers in neurodegenerative dementias. *Alzheimer's Dement* 2020;16:262–72.
 64. Abu-Rumeileh S, Capellari S, Stanzani-Maserati M, Polisch B, Martinelli P, Caroppo P, et al. The CSF neurofilament light signature in rapidly progressive neurodegenerative dementias. *Alzheimers Res Ther* 2018;10:3.
 65. Kanata E, Golanska E, Villar-Piqué A, Karsanidou A, Dafou D, Xanthopoulos K, et al. Cerebrospinal fluid neurofilament light in suspected sporadic Creutzfeldt-Jakob disease. *J Clin Neurosci* 2019;60:124–7.
 66. Steinacker P, Blennow K, Halbgebauer S, Shi S, Ruf V, Oeckl P, et al. Neurofilaments in blood and CSF for diagnosis and prediction of onset in Creutzfeldt-Jakob disease. *Sci Rep* 2016;6:38737.
 67. Thompson AGB, Luk C, Heslegrave AJ, Zetterberg H, Mead SH, Collinge J, et al. Neurofilament light chain and tau concentrations are markedly increased in the serum of patients with sporadic Creutzfeldt-Jakob disease, and tau correlates with rate of disease progression. *J Neurol Neurosurg Psychiatry* 2018;89:955–61.
 68. Gao L, Tang H, Nie K, Wang L, Zhao J, Gan R, et al. Cerebrospinal fluid alpha-synuclein as a biomarker for Parkinson's disease diagnosis: a systematic review and meta-analysis. *Int J Neurosci* 2015;125:645–54.
 69. Bousiges O, Philippi N, Lavaux T, Perret-Liaudet A, Lachmann I, Schaeffer-Agalède C, et al. Differential diagnostic value of total alpha-synuclein assay in the cerebrospinal fluid between Alzheimer's disease and dementia with Lewy bodies from the prodromal stage. *Alzheimers Res Ther* 2020;12:120.
 70. Llorens F, Kruse N, Karch A, Schmitz M, Zafar S, Gotzmann N, et al. Validation of α -synuclein as a CSF biomarker for sporadic Creutzfeldt-Jakob disease. *Mol Neurobiol* 2018;55:2249–57.
 71. Llorens F, Kruse N, Schmitz M, Gotzmann N, Golanska E, Thüne K, et al. Evaluation of α -synuclein as a novel cerebrospinal fluid biomarker in different forms of prion diseases. *Alzheimer's Dement* 2017;13:710–9.
 72. Schmitz M, Villar-Piqué A, Llorens F, Gmitterová K, Hermann P, Vargas D, et al. Cerebrospinal fluid total and phosphorylated α -synuclein in patients with Creutzfeldt-Jakob disease and synucleinopathy. *Mol Neurobiol* 2019;56:3476–83.
 73. Aksamit AJ, Preissner CM, Homburger HA. Quantitation of 14-3-3 and neuron-specific enolase proteins in CSF in Creutzfeldt-Jakob disease. *Neurology* 2001;57:728–30.
 74. Zerr I, Bodemer M, Racker S, Grosche S, Poser S, Weber T, et al. Cerebrospinal fluid concentration of neuron-specific enolase in diagnosis of Creutzfeldt-Jakob disease. *Lancet* 1995;345:1609–10.
 75. Otto M, Wiltfang J, Schutz E, Zerr I, Otto A, Pfahlberg A, et al. Diagnosis of Creutzfeldt-Jakob disease by measurement of S100 protein in serum: prospective case-control study. *BMJ* 1998;316:577–82.
 76. Vanni S, Moda F, Zattoni M, Bistaffa E, De Cecco E, Rossi M, et al. Differential overexpression of SERPINA3 in human prion diseases. *Sci Rep* 2017;7:15637.
 77. Le Pera M, Urso E, Sprovieri T, Bossio S, Aguglia U, Manna I, et al. Contribution of cerebrospinal fluid thymosin β 4 levels

- to the clinical differentiation of Creutzfeldt-Jakob disease. *Arch Neurol*. 2012;69(7):868-72.
78. Barria MA, Libori A, Mitchell G, Head MW. Susceptibility of human prion protein to conversion by chronic wasting disease prions. *Emerg Infect Dis* 2018;24:1482-9.
 79. Moda F, Gambetti P, Notari S, Concha-Marambio L, Catania M, Park K-W, et al. Prions in the urine of patients with variant Creutzfeldt-Jakob disease. *N Engl J Med* 2014;371:530-9.
 80. Concha-Marambio L, Pritzkow S, Moda F, Tagliavini F, Ironside JW, Schulz PE, et al. Detection of prions in blood from patients with variant Creutzfeldt-Jakob disease. *Sci Transl Med* 2016;8:370ra183.
 81. Bélondrade M, Nicot S, Mayran C, Bruyere-Ostells L, Almela F, Di Bari MA, et al. Sensitive protein misfolding cyclic amplification of sporadic Creutzfeldt-Jakob disease prions is strongly seed and substrate dependent. *Sci Rep* 2021;11:4058.
 82. Atarashi R, Satoh K, Sano K, Fuse T, Yamaguchi N, Ishibashi D, et al. Ultrasensitive human prion detection in cerebrospinal fluid by real-time quaking-induced conversion. *Nat Med* 2011;17:175-8.
 83. Franceschini A, Baiardi S, Hughson AG, McKenzie N, Moda F, Rossi M, et al. High diagnostic value of second generation CSF RT-QuIC across the wide spectrum of CJD prions. *Sci Rep* 2017;7:10655.
 84. Orrú CD, Bongianni M, Tonoli G, Ferrari S, Hughson AG, Groveman BR, et al. A test for Creutzfeldt-Jakob disease using nasal brushings. *N Engl J Med* 2014;371:519-29.
 85. Mammanna A, Baiardi S, Rossi M, Franceschini A, Donadio V, Capellari S, et al. Detection of prions in skin punch biopsies of Creutzfeldt-Jakob disease patients. *Ann Clin Transl Neurol* 2020;7:559-64.
 86. Cazzaniga FA, De Luca CMG, Bistaffa E, Consonni A, Legname G, Giaccone G, et al. Cell-free amplification of prions: Where do we stand? *Prog Mol Biol Transl Sci* 2020;175:325-58.
 87. Schmitz M, Cramm M, Llorens F, Müller-Cramm D, Collins S, Atarashi R, et al. The real-time quaking-induced conversion assay for detection of human prion disease and study of other protein misfolding diseases. *Nat Protoc* 2016;11:2233-42.
 88. McGuire LI, Peden AH, Orrú CD, Wilham JM, Appleford NE, Mallinson G, et al. Real time quaking-induced conversion analysis of cerebrospinal fluid in sporadic Creutzfeldt-Jakob disease. *Ann Neurol* 2012;72:278-85.
 89. Orrú C, Hughson A, Groveman B, Campbell K, Anson K, Manca M, et al. Factors that improve RT-QuIC detection of prion seeding activity. *Viruses* 2016;8:140.
 90. Candelise N, Baiardi S, Franceschini A, Rossi M, Parchi P. Towards an improved early diagnosis of neurodegenerative diseases: the emerging role of in vitro conversion assays for protein amyloids. *Acta Neuropathol Commun* 2020;8:117.
 91. Shaked GM, Fridlander G, Meiner Z, Taraboulos A, Gabizon R. Protease-resistant and detergent-insoluble prion protein is not necessarily associated with prion infectivity. *J Biol Chem* 1999;274:17981-6.
 92. McGuire LI, Poggioli A, Poggiolini I, Suardi S, Grznarova K, Shi S, et al. Cerebrospinal fluid real-time quaking-induced conversion is a robust and reliable test for sporadic creutzfeldt-jakob disease: An international study. *Ann Neurol* 2016;80:160-5.
 93. Cramm M, Schmitz M, Karch A, Mitrova E, Kuhn F, Schroeder B, et al. Stability and reproducibility underscore utility of RT-QuIC for diagnosis of Creutzfeldt-Jakob disease. *Mol Neurobiol* 2016;53:1896-904.
 94. Groveman BR, Orrú CD, Hughson AG, Bongianni M, Fiorini M, Imperiale D, et al. Extended and direct evaluation of RT-QuIC assays for Creutzfeldt-Jakob disease diagnosis. *Ann Clin Transl Neurol* 2017;4:139-44.
 95. Park J-H, Choi Y-G, Lee Y-J, Park S-J, Choi H-S, Choi K-C, et al. Real-Time Quaking-induced conversion analysis for the diagnosis of sporadic Creutzfeldt-Jakob disease in Korea. *J Clin Neurol* 2016;12:101.
 96. Bongianni M, Orrú C, Groveman BR, Sacchetto L, Fiorini M, Tonoli G, et al. Diagnosis of human prion disease using real-time quaking-induced conversion testing of olfactory mucosa and cerebrospinal fluid samples. *JAMA Neurol* 2017;74:15-62.
 97. Rhoads DD, Wrona A, Foutz A, Blevins J, Glisic K, Person M, et al. Diagnosis of prion diseases by RT-QuIC results in improved surveillance. *Neurology* 2020;95:e1017-26.
 98. Xiao K, Yang XH, Zou WQ, Dong XP, Shi Q. Assessment of the sensitivity and specificity of the established real-time quaking-induced conversion (RT-QuIC) technique in Chinese CJD surveillance. *Biomed Environ Sci* 2020;33:620-2.
 99. Orrú CD, Groveman BR, Foutz A, Bongianni M, Cardone F, McKenzie N, et al. Ring trial of 2nd generation RT-QuIC diagnostic tests for sporadic CJD. *Ann Clin Transl Neurol* 2020;7:2262-71.
 100. Fiorini M, Iselle G, Perra D, Bongianni M, Capaldi S, Sacchetto L, et al. High diagnostic accuracy of RT-QuIC assay in a prospective study of patients with suspected sCJD. *Int J Mol Sci* 2020;21:880.
 101. Hermann P, Laux M, Glatzel M, Matschke J, Knipper T, Goebel S, et al. Validation and utilization of amended diagnostic criteria in Creutzfeldt-Jakob disease surveillance. *Neurology* 2018;91:e331-8.
 102. Redaelli V, Bistaffa E, Zanusso G, Salzano G, Sacchetto L, Rossi M, et al. detection of prion seeding activity in the olfactory mucosa of patients with fatal familial insomnia. *Sci Rep* 2017;7:46269.
 103. Orrú CD, Groveman BR, Raymond LD, Hughson AG, Nonno R, Zou W, et al. Bank vole prion protein as an apparently universal substrate for RT-QuIC-based detection and discrimination of prion strains. *Supattapone S, editor. PLoS Pathog* 2015;11:e1004983.
 104. Bistaffa E, Vuong TT, Cazzaniga FA, Tran L, Salzano G, Legname G, et al. Use of different RT-QuIC substrates for detecting CWD prions in the brain of Norwegian cervids. *Sci Rep* 2019;9:18595.
 105. Orrú CD, Yuan J, Appleby BS, Li B, Li Y, Winner D, et al. Prion seeding activity and infectivity in skin samples from patients with sporadic Creutzfeldt-Jakob disease. *Sci Transl Med* 2017;9:eaam7785.
 106. Orrú CD, Soldau K, Cordano C, Llibre-Guerra J, Green AJ, Sanchez H, et al. Prion seeds distribute throughout the eyes of sporadic Creutzfeldt-Jakob disease patients. *MBio* 2018;9:e02095-18.
 107. Foutz A, Appleby BS, Hamlin C, Liu X, Yang S, Cohen Y, et al. Diagnostic and prognostic value of human prion detection in cerebrospinal fluid. *Ann Neurol* 2017;81:79-92.
 108. Piconi G, Peden AH, Barria MA, Green AJE. Epitope mapping of the protease resistant products of RT-QuIC does not allow the discrimination of sCJD subtypes. *PLoS One* 2019;14:e0218509.
 109. Hermann P, Appleby B, Brandel J-P, Caughey B, Collins S, Geschwind MD, et al. Biomarkers and diagnostic guidelines for sporadic Creutzfeldt-Jakob disease. *Lancet Neurol* 2021;20:235-46.
 110. Rudge P, Hyare H, Green A, Collinge J, Mead S. Imaging and CSF analyses effectively distinguish CJD from its mimics. *J Neurol Neurosurg Psychiatry* 2018;89:461-6.

111. Hayashi Y, Iwasaki Y, Yoshikura N, Asano T, Mimuro M, Kimura A, et al. An autopsy-verified case of steroid-responsive encephalopathy with convulsion and a false-positive result from the real-time quaking-induced conversion assay. *Prion* 2017;11:284–92.
112. Hoover CE, Davenport KA, Henderson DM, Pulscher LA, Mathiason CK, Zabel MD, et al. detection and quantification of cwd prions in fixed paraffin embedded tissues by real-time quaking-induced conversion. *Sci Rep* 2016;6:25098.
113. Green AJE. RT-QuIC: a new test for sporadic CJD. *Pract Neurol* 2019;19:49–55.
114. Cramm M, Schmitz M, Karch A, Zafar S, Varges D, Mitrova E, et al. Characteristic CSF prion seeding efficiency in humans with prion diseases. *Mol Neurobiol* 2015;51:396–405.
115. Orrú CD, Groveman BR, Hughson AG, Zanusso G, Coulthart MB, Caughey B. Rapid and sensitive RT-QuIC detection of human Creutzfeldt-Jakob disease using cerebrospinal fluid. *MBio* 2015;6:e02451-14.
116. Baiardi S, Redaelli V, Ripellino P, Rossi M, Franceschini A, Moggio M, et al. Prion-related peripheral neuropathy in sporadic Creutzfeldt-Jakob disease. *J Neurol Neurosurg Psychiatry* 2019;90:424–7.
117. Satoh K, Fuse T, Nonaka T, Dong T, Takao M, Nakagaki T, et al. Postmortem quantitative analysis of prion seeding activity in the digestive system. *Molecules* 2019;24:4601.

Non-commercial use only

Received for publication: 29 June 2021. Accepted for publication: 7 September 2021.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2021

Licensee PAGEPress, Italy

European Journal of Histochemistry 2021; 65(s1):3298

doi:10.4081/ejh.2021.3298